



EUROPEAN PATENT APPLICATION

Application number : 91402887.3

Int. Cl.⁵ : **C12N 15/00, C12N 15/85,
C12N 15/12, A07K 67/027**

Date of filing : 28.10.91

Priority : 29.10.90 US 604905

Date of publication of application :
29.07.92 Bulletin 92/31

Designated Contracting States :
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

Applicant : **INSTITUT PASTEUR**
25-28, rue du Docteur Roux
F-75724 Paris Cédex 15 (FR)

Applicant : **INSTITUT NATIONAL DE LA
SANTÉ ET DE LA RECHERCHE MÉDICALE
(INSERM)**
101, rue de Tolbiac
F-75654 Paris Cédex 13 (FR)

Inventor : **Pringault, Eric**
30, rue du Docteur Roux
F-75015 Paris (FR)
Inventor : **Robine, Sylvie**
3, rue R. Marcheron
F-92170 Vanves (FR)
Inventor : **Huet, Christian**
35, rue M. Regnier
F-75015 Paris (FR)
Inventor : **Babinet, Charles**
158, Bd. Vincent-Auriol
F-75013 Paris (FR)
Inventor : **Louvard, Charles**
23, allée de Treviso
F-92330 Sceaux (FR)

Representative : **Desaix, Anne et al**
Ernest Gutmann - Yves Plasseraud S.A., 67
bld. Haussmann
F-75008 Paris (FR)

Villin gene promoter sequence and its use in vectors, transformed mammalian cell lines, transgenic animals, and cell lines derived from the animals.

An isolated DNA sequence encoding a promoter of the human villin gene is provided. The DNA sequence can be operatively linked to a nucleotide sequence of a gene, such as an oncogene, and incorporated in cloning and expression vectors. The vectors are useful in the production of transgenic, non-human mammals as models of colorectal cancer.

Technical Field

This invention relates to a gene promoter sequence for regulating gene expression in mammalian cells. This invention also relates to vectors containing the promoter sequence, cells transformed with the vectors, transgenic animals based on the vectors, and cell lines derived from cells in the animals.

Background Art

The ability to introduce genes into the germ line of mammals is of great interest in biology. The propensity of mammalian cells to take up exogenously added DNA and to express genes included in the DNA has been known for many years. The results of gene manipulation are inherited by the offspring of these animals. All cells of these offspring inherit the introduced gene as part of their genetic make-up. Such animals are said to be transgenic.

Transgenic mammals have provided a means for studying gene regulation during embryogenesis and in differentiation, for studying the action of oncogenes, and for studying the intricate interaction of cells in the immune system. The whole animal is the ultimate assay system for manipulated genes, which direct complex biological processes.

Transgenic animals can provide a general assay for functionally dissecting DNA sequences responsible for tissue specific or developmental regulation of a variety of genes. In addition, transgenic animals provide useful vehicles for expressing recombinant proteins and for generating precise animal models of human genetic disorders.

By means of recombinant techniques, it is possible to make constructs containing oncogenes. Many oncogene constructs have been introduced into transgenic mice and have been found to elicit tumorigenic responses. For example, mice that carry the SV40 enhancer and region coding for the large T-antigen reproducibly develop tumors of the choroid plexus, which are derived from the cells lining the ventricles of the brain. Palmiter et al., *Nature* **316**:457-60 (1985). In similar experiments, the *c-myc* oncogene was fused to the LTR of mouse mammary tumor virus. In one transgenic line, females characteristically developed mammary carcinomas during the second or third pregnancy. Stewart et al., *Cell* **38**:627-37 (1984). In other experiments, the *c-myc* oncogene was driven by immunoglobulin enhancers and gave rise to malignant lymphoid tumors in transgenic mice. Adams et al., *Nature* **318**:533-8 (1985). When an oncogenic human *ras* gene was fused to an elastase promoter/enhancer construct, transgenic mice were born with pancreatic neoplasms that appeared to be due to transformation of all of the differentiating pancreatic cells. Palmiter et al., *Ann. Rev. Genet.* **20**:465-99 (1986).

For a general discussion of gene cloning and expression in animals and animal cells, see Old and Primrose, "Principles of Gene Manipulation," Blackwell Scientific Publications, London (1989), page 255 *et seq.*

Transgenic lines, which have a predisposition to specific tumors, are of great value in the investigation of the events leading to malignant transformation. It is well known that the efficacy of an anti-cancer treatment is dependent on identification of the tumor cells that are the primary cause of the disease. This is especially true in the detection of proliferation of tumor cells of the gastrointestinal tract or of renal origin. Indeed, cancers of the digestive tract are widespread, and there is a critical need for effective treatment for these diseases. The discovery of the effective treatments can be expedited by providing an animal model that will lead to carcinogenesis, which will enable the study of the efficacy, safety, and mode of action of anti-tumoral pharmaceutical agents.

In summary, there exists a need in the art for reagents and methods for providing transgenic animal models of human tumor growth and proliferation, including transgenic animal models of human tumors of the digestive tract and renal system.

Disclosure of the Invention

Accordingly, this invention aids in fulfilling these needs in the art. Specifically, this invention provides an isolated DNA sequence encoding or corresponding to a promoter of the human villin gene. The nucleotide sequence consists essentially of:

PRV.STRIDER -> Restriction Map

DNA sequence 1991 b.p. GTCGACCTGCAG ... CTCGAGGTCGAC 1 linear

Pst I **Taq I**

BSPM I Sau3A I

Taq I Mbo I

Sal I	Dpn I	Mse I
-------	-------	-------

Hinc II	Hinc II	<u>Claj</u>	<u>Hpa I</u>	Hinc II	Dde I
---------	---------	-------------	--------------	---------	-------

Acc I BspM I Alw I Hinc II Dde I Mnl I Mbo II Acc I

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

GTCGACCTGCAGGTCACCGGATCATCGATTAGTTAACCCCTAGTTGACCCCTCAGCTCATAGATGAGAAGGGAAC 80

CACGTGACGTCAGTTCCTAGTAGCTAATCAATTGGGGATTCACCTGGGGAGTCGAGTATCTACTTCTTCCCTTC

[illegible]

1	9	19	33	42	54	69	80
---	---	----	----	----	----	----	----

1	13	20	33	46	55
1	13	24	33	46	55


1	20	34	58
---	----	----	----

Year	1990	2000	2010	2020
1	10	20	30	40
2	15	25	35	45
3	20	30	40	50

	20	21
2		
3		

5
0
5
0
5
0
5
0
5
0
5
0

5
1
1
2
2
3
3
4
4
5
5



[illegible]

279
280
309
311

Nla IV
Srf I
Ncl I
Msp I
Hpa II
Bcn I
Xma I
Sma I
Sec I
Srf I
Ncl I
Bcn I
Ava I
Mnl I
Sec I
Mnl I
Rsa I
Bln I
Sec I
Srf I
EcoR II
Bln I

ATCACTGACCCGGAGCCGAGGTGAAGTAGCCGAGATTGTGCCATTGTACTTCAGCCTGGGCACAAGAGTGAA 400
TTAGTGAACCTGGGCGCTCGGCTCCCACTTCATTCGGCTTAACACGGTAACATGAAGTCGACCCTGTCTCACCTT

• | | • | • | • | •
332 340 342 372 381 381 381
332 342 332 381 381 381
332 332 332 332 336
333 333 333 333 333
333 333 333 333 333
336

5
10
15
20
25
30
35
40
45
50
55

CTCTGTCCEGAAAAATAAATAAATAAATAAATAAAGAAGGGGACAGACAGACAAGAGTGTGACGAAATGAACAGA 480
 GAGACAGGGTTTATTATTATTATTATTATTATTATTATTCTCCCTGCTCTGTTCCACTACTGCTTACTTTGCT

Hph I
 458
 Sec I
 Bsp1286 I

Hinf I Ban II

Scrf I Sec I Scrf I
 Ecor II Nla IV
 Bstn I Scrf I Ecor II
 Sau96 I Ecor II Bstn I
 Bstn I Ava II Bstn I Sec I

AAACGGTCTCCTACTTTATATACCCCAACCTGCTCTGCTGACCAAGCAATCCTGGAGCCCTGGGACTGTGATGAA 560
 TTGCCAGAGGATGAATATGGGTTGACCAAGACGACCTGCTCCTAGACCCCTCGGACCCCTGAACCATCCAACCT

507 519 530 538
 507 519 530 539
 507 522 530 539
 522 534
 522 530 539
 526 535
 535
 539

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

Nla IV
 Nla III
 Sty I
 Sec I
 Nco I
 Mnl I
 EcoN I
 Sau96 I
 Nla IV
 Ava II
 PvuM I
 EcoO109 I
 Sty I
 Sec I

566
 573
 580
 580
 580
 581
 584
 600
 601
 614
 615
 626
 626
 627
 627
 627

Nla IV
 Bsp1286 I
 Ban II
 Mae III
 Alu I
 Fnu4H I
 Bbv I
 Fnu4H I
 Bbv I
 AluN I
 Sau3A I
 Mbo I
 Dpn I
 Mnl I

643
 643
 644
 654
 665
 666
 666
 669
 694
 695
 695
 695
 718

AGGGCTCCTACAGTTACACGGCAAGCTGCTGCTAATCAGTCGAGCAGCAATGATCAGAGACTGGGAGGGCTTACCT 720
 TCCCCGAGGATGTCAATGTGCCGTCGACGAGATTAGTCAGCCTCGTCCGTTACTAGTCTCTGACCCGTCGCCGAATGA

669
649

Sec I Fnu4H I
Srf I Fnu4H I
Ecor II Bbv I
Bstn I Bbv I
Xba I
Mae I
Dde I
Mnl I
Bsu36 I
Dde I
Sca I
Rsa I
Hae III
Hae I
Mae III

727 736 744 756 763 780 783 792 799
728 736 744 756 757 784 793

736 747 757 784 793

pie I
Hinf I
Hph I
Mnl I
Dde I
Mae III

803 807 809 812 812


861 863 872 873 874

880

CCGAACCTCTAGAACGCCAGGAGCAGCAGCCACCCCTCAGGCTCAGAAAGGAGTTCTAAGTACTAAAGGCCAAGT 800
GGCTGAGATCTGCGGTCCTCGTCGTCGGTGGGAGTCCGAGCTTTTCCCTCAAGATTCATGATTTCCGGTCA

CACTAAGAGGTGACTCAAGTCAAGTCAAGTCAAGAGAGATAGATAGTGAATTCAGGCTGTCACGAGTGGAGGCTGA 880
GTGATTCCTCAGTCAAGTTCAGTCCAGTTCCTCTAATCTAATCACTCTAAAGTCCGACAGTGCCTCACCCTCCGACT

[illegible]



33

[illegible]

[illegible]

[illegible]

5
10
15
20
25
30
35
40
45
50
55

Sau96 I
Nla IV
Ava II
Mae I
Mae III
Bsp1286 I
Hae III
Mnl I
Bstn I
Ecor II
Bstn I

HolA I
Sec I
Dde I
Mnl I

AAGCGAATGTCCTGAGTGGGAGTAACCCCTGTGCTCCCCAGGGCCCTGAGCCTGGACCTCCTGGCACTGTG
TTGCCCTTACCAGGACTACCCCTCATTTGGAGACACGAGGGGTCCGGGACTCGGACCGTGGAGACCGTGACAC


1771
1771
1771
1776
1787
1792
1797
1804
1805
1805
1805
1807
1808
1808
1813
1825
1828
1828
1828

Fok I
SfaNI I
Fnu4H I
Bbv I
Dde I
Bsp1286 I
Nla IV
Bam I
Srf I
Ecor II
Bstn I
Sec I

Mae I
Fnu4H I

1849
1867
1867
1870
1871
1879
1883
1883
1884
1910
1911
1911
1911
1911

1920



This invention also provides a DNA sequence comprising the promoter of the human villin gene operatively linked to a nucleotide sequence of a structural gene or an oncogene. In specific embodiments of the invention, the structural gene encodes villin or a reporter gene, such as the gene coding for luciferase.

This invention further provides isolated RNA complementary to the promoter sequence of the human villin gene and to the other DNA sequences described herein.

In another embodiment of the invention, a eukaryotic vector is provided. The vector comprises a plasmid

or viral vector containing the promoter sequence of the human villin gene.

In addition, this invention provides mammalian cells transformed with the vectors of the invention.

Also, this invention provides transgenic animals containing the promoter of the human villin gene and cell lines cultured from cells of the transgenic animals.

5

Brief Description of the Drawings

This invention will be more fully described with reference to the drawings in which:

Fig. 1' is a representation of the structure of the isolated 5'-flanking region of the villin gene;

10 Fig. 2 depicts a partial structure of the human villin gene showing the approximate location of introns and exons;

Fig. 3 depicts the nucleotide sequence of a portion of human villin cDNA, including the location of the intervening sequences (IVS) in the 5'-first 900 base pairs (bp);

15 Fig. 4'A-B depicts the 5'-region of the nucleotide sequence of DNA encoding a promoter of the human villin gene;

Fig. 4''A-N depicts the nucleotide sequence of the villin gene promoter, including the sequence of complementary strands and restriction endonuclease sites;

Fig. 5 depicts plasmid vector pSVOA and the luciferase gene employed in one embodiment of the invention;

20 Fig. 6 is a diagrammatic representation of the procedure used to construct plasmid pPrVL of the invention; and

Fig. 7 is a diagrammatic representation of the preparation of an animal model for colorectal cancers and cell cultures derived from animal tumors.

Best Mode for Carrying Out the Invention

25

1. Gene Promoter Sequence

This invention provides an isolated DNA sequence encoding a promoter of the human villin gene. Villin is a protein with a molecular weight of about 95,000 daltons and is normally present in the villi of the digestive mucosa, more especially in intestinal villi. Villin is able to bind to actin in the presence of calcium ions.

30 Villin is usually found in brush borders, which are observed at the apex of enterocytes, particularly after they have attained their final stage of differentiation. The enterocytes line the entire intestinal lumen. Micro-villi forming the brush border are assembled at the final stage of differentiation of the enterocyte. Villin is localized in bundles of axial microfilaments of micro-villi. It contributes to the structure of their cytoskeleton. On examining frozen sections by immunofluorescence, it has been observed that villin is present at the apical pole of elongated cells forming columns at the surface of the internal wall of the intestine, and also close to cells of the proximal tubules of the kidney. In both cases, their regions are contiguous with the brush border of the cells concerned.

40 Villin has not been detected in other types of micro-villi of various other epithelia cells when the micro-villi were not provided with a well-organized brush border. A. Bretscher et al., *Exp. Cell Res.*, **135**:213 (1981); H. Reggio et al., "Membranes in Growth and Development" by A. Liss, New York (1982) pp. 89-105. Furthermore, it has been observed that villin is present in enterocytes from an early stage of their development, well before the organization of the brush border.

45 Villin can almost always be detected *in vivo* at all stages of the development of tumor cells in the digestive region, at both an early and later stages of tumorigenesis, irrespective of the state of differentiation of the cells concerned. In particular, villin can be detected in cancers of the colon, one of the most widespread forms of cancer. Villin can also be detected in certain types of renal cancer, but has never been detected in primary tumor cells derived from cancers localized in the tissues of other organs, such as the liver, ovaries, and lungs.

50 The nucleotide sequence and the amino acid sequence of the COOH-terminal portion of human villin are given below:

55

LYS TRP SER ASN THR LYS SER TYR GLU ASP LEU LYS ALA GLU
 AAG TGG AGT AAC ACC AAA TCC TAT GAG GAC CTG AAG GCG GAG
 5 SER GLY ASN SER ARG ASP TRP SER GLN ILE THR ALA GLU VAL
 TCT GGC AAC TCT AGG GAC TGG AGC CAG ATC ACT GCT GAG GTC
 THR SER PRO LYS VAL ASP VAL PHE ASN ALA ASN SER ASN LEU
 ACA AGC CCC AAA GTG GAC GTG TTC AAT GCT AAC AGC AAC CTC
 10 SER SER GLY PRO LEU PRO ILE PHE PRO LEU GLU GLN LEU VAL
 AGT TCT GGG CCT CTG CCC ATC TTC CCC CTG GAG CAG CTA GTG
 ASN LYS PRO VAL GLU GLU LEU PRO GLU GLY VAL ASP PRO SER
 15 AAC AAG GCT GTA GAG GAG CTC CCC GAG GGT GTG GAC CCC AGC
 ARG LYS GLU GLU HIS LEU SER ILE GLU ASP PHE THR GLN ALA
 AGG AAG GAG GAA CAC CTG TGC ATT GAA GAT TTC ACT GAG GCC
 PHE GLY MET THR PRO ALA ALA PHE SER ALA LEU PRO ARG TRP
 20 TIT GGG ATG ACT CCA GCT GCC TTC TCT GCT CTG CCT CGA TGG
 LYS GLN GLN ASN LEU LYS LYS GLU LYS GLY LEU PHE *
 AAG CAA CAA AAC CTC AAG AAA GAA AAA GGA CTA ITT TGA GAAG
 25 AGTAGCTGTCCTTGTAAGCAGTACCCTACCCTGATTGTAGGGTCTCATTCTCA
 CCGATATTAGTCCTACACCAATTGAAGTGAAATTTTGCAGATGTGCCTATGAGCAC
 AAACCTCTGTGGCAAATGCCAGTTTGTTTAATAAATGTACCTATTCCTTCAGAAA
 GATGATACCCCAAAAAAAAAA

30 This nucleotide sequence and amino acid sequence identify a domain, which has been termed the headpiece (HP).

The nucleotide sequence coding for the terminal portion of human villin can be isolated by means of the following procedure:

- 35 – total mRNA (messenger RNAs) is prepared according to known techniques from a cell line expressing villin;
- a cDNA (complementary DNA) bank is constructed;
- these cDNAs are inserted into a vector capable of expressing the protein coded by the insert;
- by means of the recombinant vectors obtained, a bacterial strain is transformed, then conditions permitting the expression of the desired protein in the bacteria are established; and
- 40 – the recombinant clones containing the clone specific for villin is selected by means of antibodies recognizing villin.

These techniques are described in more detail in copending U.S. application Serial No. 287,658, filed December 21, 1988. The sequence of the entire human villin protein has been elucidated by Arpin et al., J. Cell Biol., 107:1759-1766 (1988) and Pringault et al., EMBO J., 5:3119-3124 (1986).

45 A plasmid containing the gene encoding human villin was deposited with the Collection Nationale de Cultures de Microorganismes (C.N.C.M.) of Institut Pasteur in Paris, France on November 13, 1985, under culture collection deposit Accession No. I-497.

A portion of the operon encoding the human villin gene is diagrammatically shown in Fig. 1'. More particularly, Fig. 1' depicts the structure of an isolated 5'-flanking region of the operon for the human villin gene. Referring to the Figure, a 2 kb region is shown proximate the 5'-end. This region contains the gene promoter of the invention. The CAP site and the ATG start codon of the gene encoding human villin are also shown in the Figure.

It has been discovered that the operon encoding human villin contains a number of exons and introns. The first exon is shown in Fig. 1' and is identified as "EXON 1". A number of other exons and introns are depicted in Fig. 2, which is a partial structure of an isolated 5'-flanking region of the operon encoding human villin mRNA. Exon 1 in Fig. 1' corresponds to the first exon comprising nucleotides 1-97 in Fig. 2. Introns are identified by triangular regions extending toward the bottom of the Figure. It will be apparent from Fig. 2 that the operon encoding the human villin gene contains a number of intervening sequences.

The location of the intervening sequences (IVS) in a 5'-first 900 bp of human villin cDNA is shown in Fig. 3. The IVS are depicted relative to the nucleotide sequence encoding human villin. The sequence shown in Fig. 3 is the coding sequence (exons) of the villin gene. Once again, the CAP site and ATG start codon are shown in the Figure.

5 The DNA sequence of the invention encodes a promoter of the human villin gene. The DNA sequence is shown in Figs. 4'A-B and 4"A-N. Fig. 4'A-B depicts a portion of the DNA in the 5'-region of the promoter, and includes the CAP site and the ATG codon of the villin gene. Fig. 4"A-N depicts the entire 1991 bp DNA sequence encoding the promoter of the villin gene of the invention. The locations of restriction endonuclease sites in the DNA sequence are indicated in Fig. 4"A-N.

10 The nucleotide sequence of the promoter of the invention was derived by dideoxynucleotide sequencing. The base sequences of the nucleotides are written in the 5' → 3' direction. Each of the letters shown is a conventional designation for the following nucleotides:

A Adenine

G Guanine

15 T Thymine

C Cytosine.

The DNA sequence encoding the promoter of the human villin gene can be isolated from its natural environment. It is preferred that the DNA sequence encoding the promoter of the human villin gene be in a purified form. For instance, the promoter can be free of human blood-derived proteins, human serum proteins, viral proteins, nucleotide sequences encoding these proteins, human tissue, human tissue components, or combinations of these substances. In addition, it is preferred that the promoter sequence is free of extraneous proteins and lipids, and adventitious microorganisms, such as bacteria and viruses. The essentially purified and isolated DNA sequence encoding the promoter is especially useful for preparing nucleotide probes and expression vectors.

25 The promoter of the invention can also be prepared by the formation of 3' → 5' phosphate linkages between nucleoside units using conventional chemical synthesis techniques. For example, the well-known phosphodiester, phosphotriester, and phosphite triester techniques, as well as known modifications of these approaches, can be employed. Deoxyribonucleotides can be prepared with automatic synthesis machines, such as those based on the phosphoramidite approach. Oligo- and polyribonucleotides can also be obtained with the aid of RNA ligase using conventional techniques.

30 This invention of course includes variants of the DNA sequence of the invention exhibiting substantially the same properties as the promoter of the invention. By this it is meant that DNA sequences need not be identical to the sequence disclosed herein. Variations can be attributable to single or multiple base substitutions, deletions, or insertions or local mutations involving one or more nucleotides not substantially detracting from the properties of the DNA sequence as a promoter for the human villin gene. It will also be understood that the present invention is intended to encompass fragments of the DNA sequence of the invention in purified form, where the fragments are capable of functioning as promoters for the human villin gene. For example, regulatory elements responsible for tissue specificity of villin can be characterized by a functional test using the Firefly luciferase gene as a reporter gene as described by de Wet et al., *infra*.

40 The DNA sequence coding for the promoter of the present invention can be amplified in the well known polymerase chain reaction (PCR), which is useful for amplifying all or specific regions of the promoter. See e.g., S. Kwok et al., *J. Virol.*, **61**:1690-1694 (1987); U.S. Patent 4,683,202; and U.S. Patent 4,683,195. More particularly, DNA primers pairs of known sequence positioned 10-300 base pairs apart that are complementary to the plus and minus strands of the DNA to be amplified can be prepared by well known techniques for the synthesis of oligonucleotides. One end of each primer can be extended and modified to create restriction endonuclease sites when the primer is annealed to the DNA. The PCR reaction mixture can contain the DNA, the DNA primer pairs, four deoxyribonucleoside triphosphates, MgCl₂, DNA polymerase, and conventional buffers. The DNA can be amplified for a number of cycles. It is generally possible to increase the sensitivity of detection by using a multiplicity of cycles, each cycle consisting of a short period of denaturation of the DNA at an elevated temperature, cooling of the reaction mixture, and polymerization with the DNA polymerase. Amplified sequences can be detected by the use of a technique termed oligomer restriction (OR). See, R. K. Saiki et al., *Bio/Technology* **3**:1008-1012 (1985).

55 The villin gene is strictly regulated during embryonic development of the digestive tract. Villin can be detected in the visceral endoderm, in the primitive gut, in pancreatic and hepatic anlagen, as soon as these tissues appear. This expression is then stimulated in differentiated enterocytes, remains low in biliary cells and duct cells, and shut off in hepatocytes and in acinar pancreatic cells. The DNA sequence encoding the promoter of the human villin gene is thus useful to investigate molecular genetics involved in digestive organogenesis. The villin gene promoter strictly regulates the expression of villin in a tissue specific manner, during adult life and

during embryonic development.

2. Nucleotide Probes Containing the Gene Promoter

5 The DNA sequence of the invention coding for the promoter of the human villin gene can also be used as a probe for the detection of a nucleotide sequence in a biological material, such as tissue or body fluids. The probe can be labeled with an atom or inorganic radical, most commonly using a radionuclide, but also perhaps with a heavy metal. Radioactive labels including ^{32}P , ^3H , ^{14}C , or the like. Any radioactive label can be employed, which provides for an adequate signal and has sufficient half-life. Other labels include ligands that can serve
10 as a specific binding member to a labeled antibody, fluorescers, chemiluminescers, enzymes, antibodies which can serve as a specific binding pair member for a labeled ligand, and the like. The choice of the label will be governed by the effect of the label on the rate of hybridization and binding of the probe to the DNA or RNA. It will be necessary that the label provide sufficient sensitivity to detect the amount of DNA or RNA available for hybridization.

15 When the promoter sequence of the invention is used as a probe for hybridizing to a gene, the gene is preferably affixed to a water insoluble solid, porous support, such as nitrocellulose paper. Hybridization can be carried out using labeled polynucleotides of the invention and conventional hybridization reagents. The particular hybridization technique is not essential to the invention.

The amount of labeled probe present in the hybridization solution will vary widely, depending upon the nature of the label, the amount of the labeled probe which can reasonably bind to the support, and the stringency of the hybridization. Generally, substantial excesses of the probe over stoichiometric will be employed to enhance the rate of binding of the probe to the fixed DNA.

Various degrees of stringency of hybridization can be employed. The more severe the conditions, the greater the complementarity that is required for hybridization between the probe and the polynucleotide for duplex formation. Severity can be controlled by temperature, probe concentration, probe length, ionic strength, time, and the like. Conveniently, the stringency of hybridization is varied by changing the polarity of the reactant solution. Temperatures to be employed can be empirically determined or determined from well known formulas developed for this purpose.

30 3. Nucleotide Sequences Containing the Gene Promoter

This invention also provides the DNA sequence encoding the promoter of the invention linked to nucleic acids, such as nucleic acids comprising a structural gene. The nucleic acid can be obtained from any source, for example, from plasmids, from cloned DNA or RNA, or from natural DNA or RNA from any source, including
35 prokaryotic and eukaryotic organisms. DNA or RNA can be extracted from a biological material, such as biological fluids or tissue, by a variety of techniques including those described by Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York (1982). The nucleic acid will generally be obtained from a bacteria, yeast, virus, or a higher organism, such as a plant or animal. The nucleic acid can be a fraction of a more complex mixture, such as a portion of a gene contained in whole human DNA or a portion
40 of a nucleic acid sequence of a particular microorganism. The nucleic acid can be a fraction of a larger molecule or the nucleic acid can constitute an entire gene or assembly of genes. The DNA can be in a single-stranded or double-stranded form. If the fragment is in single-stranded form, it can be converted to double-stranded form using DNA polymerase according to conventional techniques.

The DNA sequence coding for the promoter of the human villin gene can be linked to a structural gene. As
45 used herein, the term "structural gene" refers to a DNA sequence that encodes through its template or messenger mRNA a sequence of amino acids characteristic of a specific protein or polypeptide. The nucleotide sequence of the invention can function as an expression control sequence, that is, a DNA sequence that controls and regulates expression of the structural gene when operatively linked to the gene. In one embodiment of this invention, the DNA sequence encoding the human villin gene promoter is operatively linked to the structural gene for human villin. In another embodiment of the invention, the DNA sequence of the invention is operatively linked to a structural gene other than the gene encoding human villin.

The nucleotide sequence of the invention coding for the promoter of the human villin gene can be used in transient assay systems based on the use of fusion genes. The fusion gene can consist of the promoter of the invention directing the synthesis of a reporter molecule. The relative amount of the reporter protein synthesized
55 under various conditions is presumed to reflect the ability of the promoter to direct transcription. The transient expression system should be based on the synthesis of an easily assayed reporter protein having minimal or no effect on the physiology of the transfected cell. Examples of suitable reporter systems are chloramphenicol acetyltransferase (CAT) and human growth hormone (hGH). Other systems that are also useful include the

luciferase system and β -galactosidase, which have been successfully used as reporter genes in eukaryotic cells. Thus, for example, the DNA sequence of the invention can be operatively linked to a reporter gene, such as the gene encoding human growth hormone (hGH) or the gene encoding for the enzyme luciferase. The Firefly luciferase gene, its structure, and its expression in mammalian cells are described in De Wet et al., Mol. Cell. Biol. 7:725-737 (1987).

The DNA sequence encoding the promoter of the human villin gene can also be linked to an oncogene. As used herein, the term "oncogene" means a gene having oncogenic or cancer causing potential. The term as used herein includes cellular oncogenes known as proto-oncogenes, as well as viral oncogenes. When the DNA sequence encoding the promoter for the human villin gene is operatively linked to an oncogene, the resulting polynucleotide is referred to herein as an "activated oncogene sequence."

Previous studies have demonstrated that villin is a marker of digestive epithelium cell lineage. Villin gene expression is strictly regulated during embryogenesis and terminal differentiation. Furthermore, the tissue specific expression of villin is maintained during the carcinogenesis process. Villin can be considered as a marker for colorectal carcinoma and can be used to characterize the colorectal origin of unknown metastasis. As a consequence of these properties, the villin gene is a good candidate to target the expression of oncogenes in the digestive epitheliums of transgenic mice. Oncogenes such as SV40T antigen, whose ability to promote tumors in transgenic mice has been fully demonstrated, K-ras or p53, whose mutated forms have been found associated with colorectal cancers in humans, can be placed under the control of the villin promoter of the invention. This makes it possible to obtain transgenic mice developing specific tumors deriving from intestinal epitheliums.

Thus, in a preferred embodiment of the invention, the DNA sequence coding for the promoter of the human villin gene is operatively linked to an oncogene that is associated with tumor cells of the gastrointestinal tract or renal system of the human. Oncogenes associated with colon carcinoma are particularly preferred for use in this invention. Examples of particularly preferred oncogenes are TAG, *ras*, *myc*, and p53 genes. It will be understood, however, that other oncogenes can be linked to the promoter sequence of the invention. For example, the promoter sequence can be linked to oncogenes that have appeared in retroviruses, such as the oncogenes *abl*, *erb-A*, *erb-B*, *ets*, *fos/fps*, *fgr*, *ms*, *fos*, *kit*, *mlf/raf*, *mos*, *myb*, *K-ras*, *rel*, *ros*, *sis*, *ski*, *src*, *yes*. Similarly, the promoter sequence of the invention can be linked to oncogenes that have not appeared in retroviruses, such as the oncogenes *bcl-1*, *bcl-2*, *bcr*, *Blym*, *dil*, *int-1*, *int-2*, *L-myc*, *mcf2*, *mcf3*, *met*, *Mlvi-1*, *Mlvi-2*, *Mlvi-3*, *neu*, *N-myc*, *N-ras*, *onc-D*, *pim-1*, *pvt-1*, *RMO-int-1*, *tel-1*, *Tlym-1*, *Tlym-2*, *TkNS-1*, *tx-1*, *tx-2*, *tx-3*, *tx-4*.

The protein products of many of the oncogenes are protein kinases or protein phosphorylating enzymes. Thus, the activated oncogene sequence can be used to study host expression mechanisms by detecting expression of these products.

4. Vectors Containing the Gene Promoter

This invention also provides cloning and expression vectors containing the DNA sequence of the invention coding for the promoter of the human villin gene.

More particularly, the DNA sequence encoding the promoter can be ligated to a vehicle for cloning the sequence. The major steps involved in gene cloning comprise procedures for separating DNA containing the gene of interest from prokaryotes or eukaryotes, cutting the resulting DNA fragment and the DNA from a cloning vehicle in specific sites, mixing the two DNA fragments together, and ligating the fragments to yield a recombinant DNA molecule. The recombinant molecule can then be transferred into a host cell, and the cells allowed to replicate to produce identical cells containing clones of the original DNA sequence.

The vehicle employed in this invention can be any double-stranded DNA molecule capable of transporting the promoter of the invention into a host cell and capable of replicating within the cell. More particularly, the vehicle must contain at least one DNA sequence that can act as the origin of replication in the host cell. In addition, the vehicle must contain two or more sites for insertion of the DNA sequence encoding the promoter of the invention. These sites will ordinarily correspond to restriction enzyme sites at which cohesive ends can be formed, and which are complementary to the cohesive ends on the promoter sequence to be ligated to the vehicle. In general, this invention can be carried out with plasmid, bacteriophage, or cosmid vehicles having these characteristics.

The nucleotide sequence of the invention can have cohesive ends compatible with any combination of sites in the vehicle. Alternatively, the sequence can have one or more blunt ends that can be ligated to corresponding blunt ends in the cloning sites of the vehicle. The nucleotide sequence to be ligated can be further processed, if desired, by successive exonuclease digestion, such as with the enzyme Bal 31. In the event that the nucleotide sequence of the invention does not contain a desired combination of cohesive ends, the sequence can be modified by adding a linker, an adaptor, or homopolymer tailing.

It is preferred that plasmids used for cloning nucleotide sequences of the invention carry one or more genes responsible for a useful characteristic, such as selectable marker, displayed by the host cell. In a preferred strategy, plasmids having genes for resistance to two different drugs are chosen. For example, insertion of the DNA sequence into a gene for an antibiotic inactivates the gene and destroys drug resistance. The second drug resistance gene is not affected when cells are transformed with the recombinants, and colonies containing the gene of interest can be selected by resistance to the second drug and susceptibility to the first drug. Preferred antibiotic markers are genes imparting chloramphenicol, ampicillin, or tetracycline resistance to the host cell.

A variety of restriction enzymes can be used to cut the vehicle. The identity of the restriction enzyme will generally depend upon the identity of the ends on the DNA sequence to be ligated and the restriction sites in the vehicle. The restriction enzyme is matched to the restriction sites in the vehicle, which in turn is matched to the ends on the nucleic acid fragment being ligated.

The ligation reaction can be set up using well known techniques and conventional reagents. Ligation is carried out with a DNA ligase that catalyzes the formation of phosphodiester bonds between adjacent 5'-phosphate and the free 3'-hydroxy groups in DNA duplexes. The DNA ligase can be derived from a variety of microorganisms. The preferred DNA ligases are enzymes from *E. coli* and bacteriophage T4. T4 DNA ligase can ligate DNA fragments with blunt or sticky ends, such as those generated by restriction enzyme digestion. *E. coli* DNA ligase can be used to catalyze the formation of phosphodiester bonds between the termini of duplex DNA molecules containing cohesive ends.

The vehicle to which the DNA sequence of the invention has been ligated is useful for cloning the sequence. Cloning can be carried in prokaryotic or eukaryotic cells. The host for replicating the cloning vehicle will of course be one that is compatible with the vehicle and in which the vehicle can replicate. When a plasmid is employed, the plasmid can be derived from bacteria or some other organism or the plasmid can be synthetically prepared. The plasmid can replicate independently of the host cell chromosome or an integrative plasmid (episome) can be employed. The plasmid can make use of the DNA replicative enzymes of the host cell in order to replicate or the plasmid can carry genes that code for the enzymes required for plasmid replication. A number of different plasmids can be employed in practicing this invention. Typical of the plasmids that can be utilized are pBR322, pBR325, ColEI, and RP4. A filamentous, lysogenic phage, such as M13 phage, can be employed for cloning the nucleotide sequences of the invention.

The DNA sequence coding for the promoter of the human villin gene and the other nucleotide sequences of the invention, such as the activated oncogene sequence, can also be ligated to a vehicle to form an expression vector. The vehicle employed in this case is one in which it is possible to express a gene operatively linked to the promoter of the invention in an appropriate host cell. It is preferable to employ a vehicle known for use in expressing genes in mammalian cells. These vehicles include SV40 vectors and eukaryotic expression vectors derived from SV40 vectors. Constructs including the DNA sequence encoding the promoter of the invention associated with the gene coding the large T antigen of SV40 are preferred, because the antigen gives transforming power to permissive cells for the expression of the promoter. Other vectors include those derived from Maloney murine sarcoma virus (MSV), herpes simplex virus, Rous sarcoma virus, the mouse mammary tumor virus (MMTV), vaccinia virus, bovine papillomavirus, human adenovirus, and Maloney murine leukemia virus.

In one embodiment of this invention, the DNA sequence coding for the promoter of the human villin gene was operatively linked to the Firefly luciferase gene, and the resulting nucleotide sequence was ligated to plasmid pSVOA using the techniques described in de Wet et al., *supra*. The plasmid pSVOA is shown in Fig. 5.

Referring to Fig. 5, the structures of the mammalian expression vector pSVOA and the full-length, intronless luciferase gene L are shown. The expression vector pSVOA is derived from pSV2, an SV40 early-region promoter vector. Portions of the vector derived from pBR322 are shown as a thin line, those from SV40 are shown as a heavy line. Coordinates of the endpoints of the pBR322 segments are indicated and correspond to the location of the restriction sites in the published sequence of pBR322. Coordinates indicated in bold type correspond to the locations of the restriction sites in the published sequence of SV40. The vector pSVOA contains the SV40 small-t-antigen intron and an SV40 polyadenylation signal labeled A_n. The vector pSVOA contains two copies of the SV40 polyadenylation signal. The full-length, intronless luciferase construct, designated L, is carried as *Hind*III-*Bam*HI fragments in the plasmid vector pUC18. The name of the corresponding plasmid is also indicated below the map. The coordinates of the indicated sites are taken from the sequence shown in Fig. 1 of the de Wet et al. publication, *supra*. Restriction sites are indicated with the following abbreviations: H3, *Hind*III; Ri, *Eco*RI; Bs, *Bsm*I; Xb, *Xba*I; H2, *Hind*II; Sp, *Ssp*I; Sa, *Sac*I; Kp, *Kpn*I; Sm, *Sma*I; Ba, *Bam*HI.

The resulting plasmid derived from pSVOA containing the DNA sequence encoding the promoter of the human villin gene operatively linked to the luciferase gene has been designated pPrVL. The plasmid pPrVL was deposited with the Collection Nationale de Cultures de Microorganismes (C.N.C.M.) of Institut Pasteur in Paris, France on October 26, 1990, under culture collection deposit Accession No. I-1011. The procedure used

to prepare the pPRVL construct is diagrammatically shown in Fig. 6.

Referring to Figure 6, details concerning the construction of pPRVL are given. A HindIII-SalI restriction fragment from pSVOAL containing the luciferase (Luc) gene was ligated to a 1.7 kb SalI-HindIII fragment obtained from a human genomic library (lambda EMBL 3). The latter fragment was obtained by screening the library using a cocktail of 5' to 3' cDNA probes specific for the human villin gene. This construct is referred to as plasmid pPrV 1.7.

A second plasmid was constructed to clone a HindIII/PstI fragment from the same library containing the CAP site of the villin gene. This plasmid was named PrV -300/+77. A HindIII-EcoRI fragment from this plasmid was then ligated to a HindIII fragment from pPrV 1.7 and an EcoRI-HindIII adaptor. The resulting plasmid (pPRVL) contained the 5' region of the villin gene (1.4 kb insert), the villin promoter and the luciferase gene.

5. Cells Transformed with Vectors Containing the Gene Promoter

The vectors of the invention can be inserted into host organisms using conventional techniques. For example, the vectors can be inserted by transformation, transfection, electroporation, microinjection, or by means of liposomes (lipofection). The preferred method for inserting the vectors into mammalian cells is dependent on cell type; however, calcium phosphate precipitation of DNA has been found to be suitable.

Cloning can be carried in prokaryotic or eukaryotic cells. The host for replicating the cloning vehicle will of course be one that is compatible with the vehicle and in which the vehicle can replicate. Cloning is preferably carried out in bacterial or yeast cells, although cells of fungal, animal, and plant origin can also be employed. The preferred host cells for conducting cloning work are bacterial cells, such as *E. coli*, as well as species of *streptomyces*, *Bacillus*, and *Pseudomonas*. The use of *E. coli* cells is particularly preferred because most cloning vehicles, such as bacterial plasmids and bacteriophages, replicate in these cells.

In a preferred embodiment of this invention, an expression vector containing the DNA sequence encoding the promoter of the invention operatively linked to a structural gene is inserted into a mammalian cell using conventional techniques. For example, the vector can be transfected into a mammalian cell by the calcium phosphate precipitation method or the DEAE-dextran method. A preferred method based on calcium phosphate precipitation is disclosed in Chen and Olsyama, *Mol. Cell. Biol.*, 7:2745-2752 (1987).

Suitable host cells include cells that express villin and cells that do not express villin. Host cells that are preferred for this purpose include human intestinal cell lines derived from colic adenocarcinoma, such as HT29 or CaCO2 cells; human epithelial cell lines, such as C33 cells; and human hepatoma cell lines, such as HepG2 cells; cell lines originating from the renal, proximal tubule of a pig, such as LLCPK1; and cells that do not express villin, such as HeLa cells, which originate from an adenocarcinoma of the neck of the uterus. After the plasmid has been introduced into the host cell, the structural gene can be expressed using conventional techniques. These cell lines can be used with recombinant vectors containing the Firefly luciferase gene under the control of the DNA sequence encoding the promoter of the invention in order to isolate regions of the promoter that can be employed for the specific expression of the villin gene in a limited number of tissues. Cell lines have been transfected with plasmid pPRVL of the invention. This plasmid contains the Firefly luciferase reporter gene.

6. Transgenic, Non-Human Mammals

Normal mice do not develop intestinal tumors spontaneously. It is possible, however, to induce adenocarcinomas from small and large intestines by using chemical agents. The histology of these tumors resembles that of corresponding human tumors, and they have been used to study external factors involved in the carcinogenesis process. The usefulness of this model is, however, limited for two main reasons. First of all, the model reproduces only some aspects of human carcinogenesis. For instance, metastasis are not observed, especially in the liver. Secondly, tumors have to be induced chemically in each mouse to be studied. This reduces the reproducibility of the model. Moreover, tumor development is very slow (average 7 months).

Transgenic mouse technology is very suitable to develop animal models of carcinogenesis, since this technology makes it possible to obtain mouse lines that genetically develop specific cancers in a reproducible manner, even those that do not appear spontaneously in this animal.

Accordingly, this invention also provides transgenic animals containing the promoter sequence of the invention. More particularly, the invention provides a transgenic non-human mammal whose germ cells and somatic cells contain an activated oncogene sequence of the invention introduced into the animal, or an ancestor of the animal (e.g. an F1 animal), at an embryonic stage. Preferably, the non-human mammal is a rodent, such as a mouse. The activated oncogene sequence, when incorporated into the genome of the mammal, increases the probability of the development of neoplasms, particularly malignant tumors, in the animal. In a

preferred embodiment of the invention, the neoplasms are analogs of human colon cancer.

There are several methods by which the oncogene sequence of the invention can be introduced into an animal embryo to ensure that it is incorporated into the chromosome of the animal in an activated state. One method is to transfect the embryo with the gene and to select transgenic animals in which the gene has integrated into the chromosome at a locus that results in activation. Other methods involve modifying the oncogene prior to introduction into the embryo. One such method is to transfect the embryo using a vector containing the activated oncogene sequence of the invention. In a preferred embodiment of the invention, the chromosome of the transgenic animal includes an endogenous coding sequence, preferably the p53, *ras*, *myc*, or *Tag* gene, which is substantially the same as the oncogene sequence, and transcription of the oncogene sequence is under the control of the promoter sequence of the invention.

Introduction of the activated oncogene sequence of the invention at the fertilized oocyte stage, such as by microinjection, ensures that the oncogene sequence will be present in all of the germ cells and somatic cells of the transgenic animal, such as the mouse. The presence of the activated oncogene sequence in the germ cells of the transgenic animal ensures that all of the descendants of the animal will carry the activated oncogene sequence in all of their germ cells and somatic cells. The activated oncogene sequence of the invention can be introduced at a later embryonic stage, although this might result in the absence of the oncogene from some somatic cells of the parent animal, although descendants of the animal that inherit the oncogene sequence will carry the sequence in all of their germ cells and somatic cells.

The animals of the invention can be used to test a material suspected of being a carcinogen by exposing the animal to the material and determining neoplastic growth as an indicator of carcinogenicity. The animals can also be used to test materials for ability to confer protection against the development of neoplasms. In either case, the animals are treated with the material, and a reduced incidence of neoplasm development compared to untreated animals is detected as an indication of carcinogenicity or protection. The invention also provides methods of exposing treated and untreated animals to carcinogen prior to, after, or simultaneously with treatment with a protective material.

The transgenic animal model of the invention provides an experimental approach to the different early and late steps of carcinogenesis, and in particular for studying oncogene implication and synergy, the apparition of proliferative hyperplasia, which occurs before the development of solid tumors, and the correlation with angiogenesis.

The villin promoter of the invention is the only promoter isolated to date, which is able to drive the expression of an oncogene in the epithelium of the large intestine. Thus, the DNA sequence encoding the promoter of the human villin gene is the only way to obtain transgenic animals developing colorectal carcinogenesis.

7. Cultures of Transgenic Animal Cells

Intestinal cell lines derived from spontaneous colorectal carcinomas are available. These cell lines have been used to establish cell culture models for absorptive enterocytes and goblet cells. However, the establishment of cell lines for PANETH entero-endocrine cells, and M cells has failed. Furthermore, no well defined cell line models for duct cells of the exocrine pancreas and for biliary cells of the liver are available.

The transgenic animals of the invention can be used as a source of cells for cell culture. Cells from the transgenic animals advantageously exhibit desirable properties of both normal and transformed cultured cells. That is, they are normal or nearly normal morphologically and physiologically, but can be cultured for longer, and even indefinite, periods of time.

More particularly, tissues of the transgenic animals of the invention can be cultured in order to obtain stable cell lines, which reproduce the differentiated phenotype of the corresponding epithelium, and which can be analyzed for the presence of the activated oncogene, either by directly analyzing DNA or RNA, or by assaying the tissue for protein expressed by the gene. Cells of tissues carrying the gene can be cultured using standard tissue culture techniques. The cells are useful for studying the functioning of cells from normally difficult to culture tissues. The cells can be used to study terminal differentiation processes. More particularly, cultures of the transgenic animal cells can be used to characterize the epithelial lines that possess specific differentiation criteria, either of the small intestine or the colon, of ductal cells of the pancreas, or of biliary cells of the liver or of the kidney proximal tubule.

Fig. 7 is a diagrammatic representation of the preparation of an animal cell model for colorectal cancers and cell cultures derived from animal tumors.

Referring to Figure 7, details are given concerning the protocol which is followed to obtain the animal and cell culture model for colorectal cancers. A vector containing the villin promoter and an oncogene (preferably p53, *ras*, *myc* or *Tag*) is transfected into a mouse oocyte. A selection procedure is used to establish which mice carry the desired genes. These animals are maintained as a colony in which the genes can be readily detected

in both somatic and germ cells.

Animals carrying the activated oncogene can then be used to investigate, for example, the induction of tumors in the digestive epithelium where villin is normally expressed. Using these animals one can develop an animal model for colorectal cancers. Alternatively, one can establish cell lines in culture from the digestive epithelium of transgenic animals carrying the villin gene and promoter which might be coupled to *ras* or *myc* to permit the *in vitro* study of colorectal tumor cells.

Claims

10

1/ An isolated DNA sequence corresponding to a promoter of the human villin gene, wherein the nucleotide sequence consists essentially of :

15

20

25

30

35

40


45

50

55

PRV.STRIDER -> Restriction Map

DNA sequence 1991 b.p. CTCGACCTGCAG ... CTGCAGCTCGAC linear



	Msp I	
	Hpa II	
	Srf I	
	Ncl I	
	Bcn I	
	Sau96 I	
	Hae III	
	Sau96 I	
	Nla IV	
	BspI286 I	
	Ban II BstU I	
	Aqa I HlnP I	
	Nla IV Hha I	
	EcoO109 I	
	Mnl I	
	Mml I	
	Alu I	
	Mae II	
	TCTACGTCATACAGCTAGTCAAGAAAAGAGGACGGCGCCGGGTGCCTCAGCCTGTAATTCCAGC	
	AGATGCAAGTATGTCGATCACCTCACAGTTCTTTCTCCTGCCCGCGCCGCCACCAGTGCGGACAATTAAGGTCG	
84	.	
94	.	
96	. .	
117	.	
124		
124		
124		132
125		132
125		133
125		
125		
125		
126		
126		
128		
128		
128		
129		
129		
160	.	



[illegible]

[illegible]

669

699

Sec I Fnu4H I Dde I Rsa I Hae III
Srf I Fnu4H I EcoR II Bbv I Mnl I Dde I Sca I Hae I Mae III
Xba I Bstn I Bbv I Bsu36 I Dde I Hae I Mae III

CCGAAGCTCTAGAACGCCAGGAGCAGCAGCAGCCACCCCTCAGGCTCAGAAAGGAGTTCTAAGTACTAAAGGCCAAGT 800
GGCTTGAGATCTTGGCGTCCCTTCGTCGTCGGTGGGAGTCCGAGCTTTTCCCTCAAGATTCATGATTTCCGGTTCA


727 736 744 756 780 792 799
728 736 747 756 783 793
736 744 757 784

Ple I Hph I Hinf I Mnl I Dra III Stu I Hae III
Mnl I Mae III Mnl I Mae III Mnl I Hae I Mnl I

CACTAAGAGGTGACTCAAGTCAAGGTGCAAGAGAGATTAGATAGTGAAGTTTCAGGCTGTCAAGGAGTGAAGGCTGA 880
GTGATTCTCCACGTGAGTTTCAGTTCCACGTTCTCTCTAATCTATCACTCTAAGTCCGACAGTGCCTCAGCTCCGAGT

803 810 807 809 812 812 812

[illegible]



35

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

Mse I BstX I Mnl I Mae III Dpn I Mnl I
 1289 1295 1307 1313 1322 1323 1323 1331 1353 1353 1353 1356 1356

ATGCGTTGTTAATCCAGCACTTGGAGGCTGTGACAGGTGATCAGTTGAGTCAAGAGTTTGAGACCAGCCCTGGCC 1360
 TACGGAACAATTAGGTCGTGAACCCCTCGACACTGTCCAGTCACTAGTCACTCAGTCCAGTCCCAAACTCTGTCGACCGG

Nla III Bcl I Mae III Alu I Mnl I
 1362 1400 1412 1429 1439

ACATGGGGAACCCCTGTCTCTACTAAATAACAATTAGCCAGCGTGGCTGTACAGCCCTGTAATCCAGCTATGCGGA 1440
 TGTACCCCTTGGACAGAGATGATTTTATGTTTAAATCGGTGCGACCGAGTGGGACATTAGGTCGATACGCCCT

Mnl I Hinf I BstX I Mnl I
 1445 1454 1466 1466 1466 1466 1471 1477 1495 1495 1495

GGTGAGGTAGGAGATCAGTTGAACCTGGAGGAGAGGTTGACAGTGAGCCAAAGATCAGACACTGCACTCCAGCCTGT 1520
 CCAACTCCATCCTCTTAGTGAACCTGACCCCTCGTCCCAACGTCACCTGGTCTAGTCTGTACGTCAGGTGCGACA

Sec I Scr I Ecor II Mnl I Sau3A I Mbo I Dpn I
 1471

5
10
15
20
25
30
35
40
45
50
55

Ple I Mbo II Nml I Alu I
Hinf I Mbo II Nla III Hind III
1533 1549 1586 1597
1533 1552 1589 1598
1555

Nla IV
Mnl I Mnl I Fok I
TTGGGAGGGGGTTCCTCCAGTTACCGGGCAGAGAGTAGTTCACAGTGGGAGGGTTCACCTGTTCATTCAATC 1680
AACCCCTCCCCCAAGAAGGTCAATCGCCCGTCTCTCATCAAGTGTCACCCTCCCAAGTGAGACAAGTAAGTAG
1606 1653 1677
1612

Mnl I Mbo II Mae III Mae I Rsa I
CATCTCTCACTCCTTCCCCTCCCCCTCTCTTCTTATGAGCAAAGTCACTGTGCTAGACACTGCCGATTGTGTAATAAC 1760
GTAGAGAGTGAGGAAGGAGGGGAGAGAAAGAAATACTCGTTTCAGTGACACGATCTGTGACCGCTAACCATGATTTG
1697 1707 1725 1734 1752
1703

Sau96 I
Nla IV
EcoO109 I
Dde I ScrF I
Ecor I ScrF I

5
10
15
20
25
30
35
40
45
50
55

Sau96 I
Nla IV
Ava II
Mae I
Mae III
Bsp1286 I
Hae III
Mnl I
Bstn I
Ecor II
Bstn I

1771
1771
1771
1776
1787
1792
1797
1804
1805
1805
1805
1807
1808
1808
1813
1825
1828
1828
1828


Fok I
SfaNI I
Fnu4H I
Bbv I
Dde I
Bsp1286 I
Nla IV
Bam I
Sbf I
Ecor II
Bstn I

Mae I
Fnu4H I
Sec I

1849
1867
1867
1870
1871
1879
1883
1883
1884
1910
1911
1911
1911

1920

1840



2/ A DNA sequence comprising the sequence as claimed in claim 1 operatively linked to a nucleotide sequence of an oncogene.

3/ The DNA sequence as claimed in claim 1 or 2, wherein the oncogene is selected from the group consisting of p53, ras, myc and Tag.

4/ An isolated DNA sequence as claimed in claim 1 operatively linked to a structural gene.

5/ An isolated DNA sequence comprising the sequence as claimed in claim 1 or 4 operatively linked to a

gene encoding villin.

6/ A DNA sequence comprising the sequence as claimed in claim 1 operatively linked to a gene encoding luciferase.

7/ An isolated RNA sequence complementary to the nucleotide sequence according to anyone of claims 1 to 6.

8/ A vehicle comprising a plasmid or viral vector containing the nucleotide sequence as claimed in anyone of claims 1 to 6.

9/ The vehicle as claimed in claim 8, wherein the viral vector is an SV-40 vector.

10/ The vehicle as claimed in claim 8, wherein the plasmid is pSVOAL.

11/ The vehicle as claimed in claim 8 having the identifying characteristics of the vector having culture collection accession number CNCM (Collection Nationale des Microorganismes, Paris, France) I-1011.

12/ The vehicle as claimed in claim 8, wherein the vector is an expression vector.

13/ A transgenic, non-human mammal all of whose germ cells and somatic cells contain a recombinant activated oncogene sequence according to anyone of claims 2 to 5, which has been introduced into the mammal, or an ancestor of the mammal, at an embryonic stage.

14/ The mammal as claimed in claim 13, wherein a chromosome of the mammal includes an endogenous coding sequence substantially the same as a coding sequence of said oncogene sequence.

15/ The mammal as claimed in claim 14, wherein the oncogene sequence has been integrated into a chromosome of the mammal at a site different from the location of the endogenous coding sequence.

16/ The mammal as claimed in anyone of claims 13 to 15, wherein the mammal is a rodent.

17/ The animal as claimed in anyone of claims 13 to 15, wherein the mammal is a mouse.

STRUCTURE OF THE ISOLATED 5'-FLANKING REGION OF THE VILLIN GENE




FIG. 1'




FIG. 2

/Cap site */Met.....*
 CCTCTCCCCCAGGCTCACTCACC ATG ACC AAG CTG AGC GCC CAA GTC AAA GGC TCT CTC
 AAC ATC ACC ACC CCG GGG CTG CAG ATA TGG AGG ATC GAG(+97) --- *IVS 1*
(390 bp) --- GCC ATG CAG ATG GTG CCT GTT CCT TCC AGC ACC TTT GGA AGC
 TTC TTC GAT GGT GAC TGC TAC ATC ATC CTG GCT(+172) --- *IVS 2 (1300*
bp) --- ATC CAC AAG ACA GCC AGC AGC CTG TCC TAT GAC ATC CAC TAC TGG
 ATT GGC CAG GAC TCA TCC CTG GAT GAG CAG GGG GCA GCT GCC ATC TAC ACC ACA
 CAG ATG GAT GAC TTC CTG AAG GGC CGG GCT GTG CAG CAC CGC GAG GTC CAG GGC
 AAC GAG AGC GAG GCC TTC CGA GGC TAC TTC AAG CAA GGC CTT GTG(+370) ---
IVS 3 (>2000 bp) --- ATC CGG AAA GGG GGC GTG GCT TCT GGC ATG AAG
 CAC GTG GAG ACC AAC TCC TAT GAC GTC CAG AGG CTG CTG CAT GTC AAG GGC AAG
 AGG AAC GTG GTA GCT GGA G(+476) --- *IVS 4 (450 bp)* --- AG GTA GAG
 ATG TCC TGG AAG AGT TTC AAC CGA GGG GAT GTT TTC CTC CTG GAC CTT GGG AAG
 CTT ATC ATC CAG TGG AAT GGA CCG GAA AGC ACC CGT ATG GAG AGA CTC AGG
 G(+590) --- *IVS 5 (1000 bp)* --- GC ATG ACT CTG GCC AAG GAG ATC
 CGA GAC CAG GAG CGG GGA GGG CGC ACC TAT GTA GGC GTG GTG GAC GGA GAG AAT
 GAA TTG GCA TCC CCG AAG CTG ATG GAG GTG ATG AAC CAC GTG CTG GGC AAG CGC
 AGG GAG CTG AAG GCG GCC GTG CCC GAC ACG GTG GTG GAG CCG GCA CTC AAG GCT
 GCA CTC AAA CTG TAC CA(+792) --- *IVS 6* --- T GTG TCT GAC TCC GAG GGG
 AAT CTG GTG GTG AGG GAA GTC GCC ACA CGG CCA CTG ACA CAG GAC CTG CTC AGT
 CAC GAG GA(+873) --- *IVS 7* --- C TGT TAC ATC

FIG. 3

GTCGA CCTGCA GGTCAA CGG ATCATCGATT TAGTTAACCC CCTAAGTTGACCCCTCAGC - 1850
 TCATAGATGAAGAAGGAAG TCTACGTT CATACAGCTAGT GAGTGTCAAAGAAAAAGAGG - 1790
 GACGGGGCCCGCGCGGTGG CTCACGCCTGTAATTCCAGC AATTTAGGAGGCTGAGGCGG - 1730
 GTGGATCACCTGAGGT CAGG AGTTCGAGACCAGCCTGGCC AACATGTGGTGGAAATCCCG - 1670
 TCTCTATTAATAAATAACAAA TTAGCTGGGCTTGAGGCTCG CGCCTGT AATCCAGCTACTT - 1610
 CAGGGAGGCTGAGGCAGGAG AATCACTTGAACCCGGGAGC CGAGGTTGAAGTAAGCCGAG - 1550
 ATTGTGCCATTGTACTTCAG CCTGGGCAACAAGAGTGAAA CTCTGTCCCAAAAATAAATA - 1490
 AATAAATAAATAAATAAAAG AAGGGGACAGAGACAAAGGT GATGACAGAAATGAAACAGA - 1430
 AAACGGTCTCCTACTTTATA CCCCAACCTGGTTCTGCTGG ACCAGGAATCCTGGGAGCCC - 1370
 TGGGGACTTGGTAGGTTGAA AAATACCTGATGGAGGAGAC CATGGTTXXXCCTTGATGG - 1310
 AGAAGGCCATAGCCAACAGG CCATCAGACAGGACCCAAGC CCCAAGGGGCTCCTACAGTT - 1250
 ACACGGCAAGCTGCTGCTAA TCAGTCGGAGCAGGCAATGA TCAGAGACTGGGCAGGGCTT - 1190
 ACCTCCGAACCTAGAACGC CAGGGAAGCAGCAGCCACCC CTCAGGCTCAGAAAAGGGAG - 1130
 TTTCTAAGTACTAAAAGGCC AAGTCACTAAGAGGTGACTC AAAGTCAAGGTGCAAGGAGA - 1070
 GATTAGATAGTGAGATTTCA GGCTGTACGGAGTGGAGGC CTGACCTGT CGGCACATGCC - 1010
 CTCTCATT CATTATTCATCA GTTCAAGGAGCATT CATGGA GGGCAGCTCTCTCGAGGCTG - 950
 GGTGATAGGCTCCATGGAGA TCGGGGTGGCCTCTCTGGAT GAGGAGGAACCTCAAACTA - 890
 GGGAGGAGAATAACTGGGCA CTAGTGGCTGTCTGCCTAGT TCATATCAGAATCACTGGGG - 830
 AGTTTTAAAAGCACCCCATG CCTCACCTCCCAAAAGGCAG AAAATAAAGCACCCATGCC - 770
 AAGCCCCACCCTGGCCTAAT TAAATCAGAATACCAGAGCC TGGAGCATCAGCACGCTXCC - 710
 GAAGCTCCCCAGCTGAGGCT GATGAGCAGCAGGAGCGAAT ACAGCTACAGGTCCAAGCAA - 650
 AAAGGGCCCGCCAGACACGT GGCTCATGCCTTGTTAATCC CAGCACTTTGGGAGGCTGTG - 590
 ACAGGTGGATCACTTGAGGT CAGGAGTTTGAGACCAGCCT GGCCACATGGGGAAACCCT - 530
 GTCTCTACTAAAAATAACAAA ATTAGCCAGCGTGGCTGTCA CGCCTGT AATCCCAGCTATG - 470
 CGGGAGGTTGAGGTAGGAGA ATCACTTGAACCTGGGAGGC AGAGGTTGCAGTGAGCCAAG - 410

FIG. 4'A

ATCACGACACTGCACTCCAG CCTGTGCGACAGAGCGGGAC TCCATCTCAAAAAGAAGAAG - 350
 AAGAAAAGAATAATATGGAG AAAGGGCTGACATGAGGAGA AAAGCTTGGGGAGGGGGGTT - 290
 CCTTCCAGTTTAGCGGGCAG AGAGTAGTTCACAGTGGGAG GGTTCACCTCTGTTTCATTCAT - 230
 TCATCCATCTCTCACTCCTT CCCTCCCCCTCTCTTCCTTA TTGAGCAAAGTCACTGTGCT - 170
 AGACACTGCGGATTTGGTAC TAAACAAGCGGAAATGGTCC CTGAGTGGGGAGTAACCCCTC - 110
 TGTGCTCCCCCAGGGCCCTG AGCCTGCGACCTCCTGGCAC TGTGGGAGTTCTCTAGGGG - 50
 CAGGCTGAAGGGCAGCATCC GAACTCAGGTGCCCTGCTC CCAACGCCT T ^{+1 (CAP site)} TCCCCCA
 GGCTCACTCACC ⁺²¹ ATG ACC AAG CTG AGC GCC CAA GTC AAA GGC TCT CTC AAC AT
 C ACC ACC CCG GGG CTG CAG ⁺⁸⁰ GTCGAC

FIG. 4'B

PRV.STRIDER -> Restriction Map

DNA sequence 1991 b.p. CTCGACCTGCAG ... CTGCAGCTCGAC 11near




FIG. 4"A

FIG. 4"B

FIG. 4"D




FIG. 4"E

[illegible]




FIG. 4"G










FIG. 4"J



[illegible]



Sec I
Scrf I
Ncl I
Msp I
Hpa II
Bcn I

59





FIG. 5

psVOAL

FIG. 6






FIG. 7



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 91 40 2887

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
	No relevant documents disclosed -----		C12N15/00 C12N15/85 C12N15/12 A07K67/027
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C12N C07K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24 FEBRUARY 1992	Examiner CHAMBONNET F. J.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>..... A : member of the same patent family, corresponding document</p>			

EPO FORM 150 (11.82) (P0401)